

ON THE VECTOR VALUED FOURIER TRANSFORM AND COMPATIBILITY OF OPERATORS*

IN SOOK PARK

ABSTRACT. Let $\mathbb{G}$ be a locally compact abelian group and let $1 < p \leq 2$. $\mathbb{G}'$ is the dual group of $\mathbb{G}$, and p' the conjugate exponent of p . An operator T between Banach spaces X and Y is said to be compatible with the Fourier transform $F^{\mathbb{G}}$ if $F^{\mathbb{G}} \otimes T : L_p(\mathbb{G}) \otimes X \rightarrow L_{p'}(\mathbb{G}') \otimes Y$ admits a continuous extension $[F^{\mathbb{G}}, T] : [L_p(\mathbb{G}), X] \rightarrow [L_{p'}(\mathbb{G}'), Y]$. $\mathcal{FT}_p^{\mathbb{G}}$ denotes the set of such T 's. We show that $\mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z}^n \times \mathbb{G}}$ for any $\mathbb{G}$ and positive integer n . And if the factor group of $\mathbb{G}$ with respect to its component of the identity element is a direct sum of a torsion free group and a finite group with discrete topology then $\mathcal{FT}_p^{\mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z}}$.

1. INTRODUCTION

A locally compact abelian group means a topological abelian group with its Haar measure whose topology is locally compact Hausdorff. The real line $\mathbb{R}$, the discrete group of integers $\mathbb{Z}$ and the circle group $\mathbb{T}$ are important examples. Further information can be found in [3], [5] and [12]. Let $\mathbb{G}$ be a locally compact abelian group and $\mathbb{G}'$ its dual(character) group, the Haar measure of $\mathbb{G}'$ is determined so that Parseval's identity is established with constant 1. For $1 \leq r < \infty$, we denote by $[L_r(\mathbb{G}, \mu_{\mathbb{G}}), X]$ the Banach space of all measurable functions $\mathbf{f} : \mathbb{G} \rightarrow X$ for which $\|\mathbf{f}\|_{[L_r(\mathbb{G}), X]} := \left(\int_{\mathbb{G}} \|\mathbf{f}(s)\|^r d\mu_{\mathbb{G}}(s) \right)^{1/r}$ is finite. Let $\mathbb{G}$ be a fixed infinite locally compact abelian group and $1 < p \leq 2$. $F^{\mathbb{G}}$ denotes the Fourier transform from $L_p(\mathbb{G})$ into $L_{p'}(\mathbb{G}')$. For a bounded linear operator T between Banach spaces X and Y ,

$$F^{\mathbb{G}} \otimes T : \sum_{k=1}^n f_k \otimes \mathbf{x}_k \mapsto \sum_{k=1}^n F^{\mathbb{G}} f_k \otimes T \mathbf{x}_k$$

yields a well-defined map from $L_p(\mathbb{G}) \otimes X$ into $L_{p'}(\mathbb{G}') \otimes Y$. T is said to be compatible with $F^{\mathbb{G}}$ or have $\mathbb{G}$ -Fourier type p , if the operator $F^{\mathbb{G}} \otimes T : L_p(\mathbb{G}) \otimes$

1991 *Mathematics Subject Classification*. Primary 46B20; Secondary 46B07.

Key words and phrases. Banach space, operator, Fourier transform, vector valued function, locally compact abelian group, dual group.

*This work is supported by BK21 project.

$X \rightarrow L_{p'}(\mathbb{G}') \otimes Y$ admits a continuous extension

$$[F^{\mathbb{G}}, T] : [L_p(\mathbb{G}), X] \rightarrow [L_{p'}(\mathbb{G}'), Y].$$

For such T , we let

$$\|T|_{\mathcal{FT}_p^{\mathbb{G}}}\| := \|[F^{\mathbb{G}}, T] : [L_p(\mathbb{G}), X] \rightarrow [L_{p'}(\mathbb{G}'), Y]\|.$$

The norm defined above is invariant under changing the Haar measure of $\mathbb{G}$. The class of these operators is a Banach ideal, denoted by $\mathcal{FT}_p^{\mathbb{G}}$. The definition and notation follow those of [10].

It is known that $\mathcal{FT}_p^{\mathbb{R}} = \mathcal{FT}_p^{\mathbb{Z}} = \mathcal{FT}_p^{\mathbb{T}}$ [10], but the problem whether the operator ideal $\mathcal{FT}_p^{\mathbb{G}}$ depends on $\mathbb{G}$ or not is unsolved. There are several results about $\text{FT}_p^{\mathbb{G}}$, the class of Banach spaces whose identity operators are compatible with $F^{\mathbb{G}}$. And these are immediately extended to the case of $\mathcal{FT}_p^{\mathbb{G}}$ by replacing the identity operator on a Banach space X with T . Peetre [9] who introduced the concept of Banach space of Fourier type p proved that X belongs to $\text{FT}_p^{\mathbb{R}}$ if and only if the dual space X' belongs to $\text{FT}_p^{\mathbb{R}}$. In fact, T belongs to $\mathcal{FT}_p^{\mathbb{G}}$ if and only if the dual operator T' belongs to $\mathcal{FT}_p^{\mathbb{G}'}$ i.e. $\|T|_{\mathcal{FT}_p^{\mathbb{G}}}\| = \|T'|_{\mathcal{FT}_p^{\mathbb{G}'}}\|$ for any locally compact abelian group $\mathbb{G}$. Bourgain [2] showed that $\text{FT}_p^{\mathbb{T}} \subset \text{FT}_p^{\mathbb{R}}$ and König [7] modified Kwapien's argument [8] to show that $\text{FT}_p^{\mathbb{R}} = \text{FT}_p^{\mathbb{T}}$ and extended this to that $\text{FT}_p^{\mathbb{G}} = \text{FT}_p^{\mathbb{T}}$ if $\mathbb{G}$ is one of $\mathbb{R}^m$ and $\mathbb{T}^m$, where m is a positive integer. Garcia-Cuerva, Kazarian and Torrea [4] and Andersson [1] showed independently that $\text{FT}_p^{\mathbb{G}} = \text{FT}_p^{\mathbb{Z}}$ whenever $\mathbb{G}$ is one of $\mathbb{T}^m$, $\mathbb{T}^{\infty}$, $\mathbb{R}^m$, $\mathbb{Z}^m$ and $\mathbb{Z}^{\infty}$. Andersson [1] also proved that $\|I_X|_{\mathcal{FT}_p^{\mathbb{H}}}\| \leq \|I_X|_{\mathcal{FT}_p^{\mathbb{G}}}\|$ when $\mathbb{H}$ is an open subgroup of $\mathbb{G}$, and that $\text{FT}_p^{\mathbb{E}} = \text{FT}_p^{\mathbb{Z}}$ if $\mathbb{E}$ is a nontrivial torsion free abelian group with the discrete topology.

In this paper we characterize $\mathcal{FT}_p^{\mathbb{G}}$ partly as follows. In Section 2, we show for every locally compact abelian group $\mathbb{G}$ that $\|T|_{\mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}}}\|$ is equivalent to $\|T|_{\mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}}\|$, and $\|T|_{\mathcal{FT}_p^{\mathbb{Z}^n \times \mathbb{G}}}\| = \|T|_{\mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}}\|$. The Cartesian product means the direct sum. By applying these, we easily obtain some results in the above paragraph and the relation $\mathcal{FT}_p^{\mathbb{R}^k \times \mathbb{Z}^l \times \mathbb{T}^m \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}$ for any nonnegative integers k, l, m with $k + l + m \geq 1$. In Section 3, we combine the results of Section 2 with the properties of a locally compact abelian group to show that $\mathcal{FT}_p^{\mathbb{R}^k \times \mathbb{Z}^l \times \mathbb{F}} = \mathcal{FT}_p^{\mathbb{Z}}$ for any compact abelian group $\mathbb{F}$ with finitely many components. And we show that if $\mathbb{G} \cong \mathbb{R}^k \times [\text{torsion-free group with discrete topology}] \times [\text{compact group with finite components}]$ then $\mathcal{FT}_p^{\mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z}}$. If the factor group of a locally compact abelian group $\mathbb{G}$ with respect to the component of the identity element is in the form of $[\text{torsion-free group}] \times [\text{finite group}]$ with discrete topology, then we have $\mathcal{FT}_p^{\mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z}}$.

From now on, X and Y are Banach spaces and $T : X \rightarrow Y$ is a bounded linear operator. We denote the dual group of $\mathbb{G}$ by $\mathbb{G}'$. But we use the fact that

$\mathbb{R}' = \mathbb{R}$ and $\mathbb{Z}' = \mathbb{T}$. We use the abbreviation LCA for ‘locally compact abelian’. The term ‘isomorphic’ means ‘topologically and algebraically isomorphic’. The integral for a vector valued function is the Bochner integral.

2. CLASSIFYING $\mathcal{FT}_p^{\mathbb{G}}$ VIA DIRECT SUM.

First we observe that the proof in [4] of the fact $\mathcal{FT}_p^{\mathbb{R}} = \mathcal{FT}_p^{\mathbb{Z}}$ can be modified to yield the following:

Proposition 1. *For any LCA group $\mathbb{G}$, we have the inequalities*

$$\|T|\mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}}\| \leq \|T|\mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}\| \leq \frac{\pi}{2} \|T|\mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}}\|,$$

and hence

$$\mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}.$$

Proof. For arbitrary $\delta > 0$, let

$$\mathbf{f}(s, t) = \sum_m \chi_{[\delta(m-\frac{1}{2}), \delta(m+\frac{1}{2})]}(s) \mathbf{g}_m(t),$$

where the summation is over $\mathbb{Z}$, $\mathbf{g}_m$ is an X -valued simple function on $\mathbb{G}$ and $\mathbf{g}_m = 0$ except for finitely many m . Note that the set of all X -valued functions of the form of $\mathbf{f}$ is dense in $[L_p(\mathbb{R} \times \mathbb{G}), X]$. We compute

$$\begin{aligned} \|\mathbf{f}|_{L_p(\mathbb{R} \times \mathbb{G})}\| &= \left(\int_{\mathbb{R} \times \mathbb{G}} \left\| \sum_m \chi_{[\delta(m-\frac{1}{2}), \delta(m+\frac{1}{2})]}(s) \mathbf{g}_m(t) \right\|^p ds dt \right)^{\frac{1}{p}} \\ (1) \quad &= \delta^{\frac{1}{p}} \left(\int_{\mathbb{G}} \sum_m \|\mathbf{g}_m(t)\|^p dt \right)^{\frac{1}{p}}. \end{aligned}$$

Since $\int_{\delta(m-1/2)}^{\delta(m+1/2)} \exp(i\tilde{s}s) ds = \exp(im\delta\tilde{s}) \frac{\sin(\frac{\delta}{2}\tilde{s})}{\frac{\delta}{2}\tilde{s}}$, we have

$$\begin{aligned} \widehat{T\mathbf{f}}(\tilde{s}, \tilde{t}) &= \int_{\mathbb{G}} \int_{\mathbb{R}} \sum_m T\mathbf{g}_m(t) \chi_{[\delta(m-\frac{1}{2}), \delta(m+\frac{1}{2})]}(s) \exp(i\tilde{s}s)(t, \tilde{t}) ds dt \\ &= \int_{\mathbb{G}} \sum_m T\mathbf{g}_m(t)(t, \tilde{t}) \exp(im\delta\tilde{s}) \frac{\sin(\frac{\delta}{2}\tilde{s})}{\frac{\delta}{2}\tilde{s}} dt. \end{aligned}$$

Hence

$$\begin{aligned}
\| [F^{\mathbb{R} \times \mathbb{G}}, T] \mathbf{f} \|^{p'} &= \int_{\mathbb{G}'} \int_{\mathbb{R}} \left\| \int_{\mathbb{G}} \sum_m T \mathbf{g}_m(t) (t, \tilde{t}) \exp(i\delta m \tilde{s}) \frac{\sin(\frac{\delta}{2} \tilde{s})}{\frac{1}{2} \tilde{s}} dt \right\|^{p'} \frac{1}{2\pi} d\tilde{s} d\tilde{t} \\
&= \delta^{p'} \int_{\mathbb{G}'} \int_{\mathbb{R}} \left\| \int_{\mathbb{G}} \sum_m T \mathbf{g}_m(t) (t, \tilde{t}) \exp(i\delta m \tilde{s}) \frac{\sin(\frac{\delta}{2} \tilde{s})}{\frac{\delta}{2} \tilde{s}} dt \right\|^{p'} \frac{1}{2\pi} d\tilde{s} d\tilde{t} \\
&= \delta^{p'-1} \int_{\mathbb{G}'} \int_{\mathbb{R}} \left| \frac{\sin(\frac{1}{2} \tilde{s})}{\frac{1}{2} \tilde{s}} \right|^{p'} \left\| \int_{\mathbb{G}} \sum_k \left(\sum_m T \mathbf{g}_m(t) \chi_{\{m\}}(k) \right) (t, \tilde{t}) \exp(ik\tilde{s}) dt \right\| \frac{1}{2\pi} d\tilde{s} d\tilde{t} \\
&= \delta^{p'-1} \int_{\mathbb{G}'} \int_{-\pi}^{\pi} \sum_n \left| \frac{\sin(\frac{1}{2} \tilde{s})}{\frac{1}{2} \tilde{s} - n\pi} \right|^{p'} \| [F^{\mathbb{Z} \times \mathbb{G}}, T] \left(\sum_m \mathbf{g}_m \chi_{\{m\}} \right) \|^{p'} \frac{1}{2\pi} d\tilde{s} d\tilde{t} \\
&\leq \delta^{p'-1} \| T | \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} \|^{p'} \left(\int_{\mathbb{G}} \sum_m \| \mathbf{g}_m(t) \|^p dt \right)^{\frac{p'}{p}} \\
&= \| T | \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} \|^{p'} \| \mathbf{f} | L_p(\mathbb{R} \times \mathbb{G}) \|^{p'} \quad \text{by (1),}
\end{aligned}$$

where we used the inequality $\sum_n \left| \frac{\sin \tilde{s}}{\tilde{s} - n\pi} \right|^{p'} \leq 1$ for any real $\tilde{s} \neq n\pi$ and $2 \leq p' < \infty$, see [6].

Therefore we have $\| T | \mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} \| \leq \| T | \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} \|$ and $\mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} \subseteq \mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}}$.

For the right inequality let $\mathbf{f}(k, t) = \sum_m \mathbf{g}_m(t) \chi_{\{m\}}(k)$ where $\mathbf{g}_m$ is an X -valued simple function and $\mathbf{g}_m = 0$ except for finitely many m . By density argument it's enough to consider $\mathbf{f}$ of the above form. Now we have

$$\begin{aligned}
\| \mathbf{f} | L_p(\mathbb{Z} \times \mathbb{G}) \| &= \left(\int_{\mathbb{G}} \sum_k \left\| \sum_m \mathbf{g}_m(t) \chi_{\{m\}}(k) \right\|^p dt \right)^{\frac{1}{p}} \\
(2) \quad &= \left(\int_{\mathbb{G}} \sum_m \| \mathbf{g}_m(t) \|^p dt \right)^{\frac{1}{p}}; \quad \text{and}
\end{aligned}$$

$$\begin{aligned}
\widehat{T\mathbf{f}}(\tilde{s}, \tilde{t}) &= \int_{\mathbb{G}} \sum_k \left(\sum_m T \mathbf{g}_m(t) \chi_{\{m\}}(k) \right) \exp(i\tilde{s}k) (t, \tilde{t}) dt \\
&= \int_{\mathbb{G}} \sum_m T \mathbf{g}_m(t) \exp(i\tilde{s}m) (t, \tilde{t}) dt.
\end{aligned}$$

We use the identity, $\exp(im\tilde{s}) = \int_{\mathbb{R}} \frac{\tilde{s}/2}{\sin(\tilde{s}/2)} \chi_{[m-1/2, m+1/2]}(s) \exp(i\tilde{s}s) ds$, to obtain the following inequality:

$$\begin{aligned}
& \| [F^{\mathbb{Z} \times \mathbb{G}}, T] \mathbf{f} \|^{p'} \\
&= \frac{1}{2\pi} \int_{\mathbb{G}'} \int_{-\pi}^{\pi} \left\| \int_{\mathbb{G}} \sum_m T \mathbf{g}_m(t) (t, \tilde{t}) \int_{\mathbb{R}} \frac{\tilde{s}/2}{\sin(\tilde{s}/2)} \chi_{[m-1/2, m+1/2]}(s) \exp(i\tilde{s}s) ds dt \right\|^{p'} d\tilde{s} d\tilde{t} \\
&\leq \frac{1}{2\pi} \int_{\mathbb{G}'} \int_{-\pi}^{\pi} \left| \frac{\tilde{s}/2}{\sin(\tilde{s}/2)} \right|^{p'} \left\| \int_{\mathbb{G}} \sum_m T \mathbf{g}_m(t) \chi_{[m-\frac{1}{2}, m+\frac{1}{2}]}(s) (t, \tilde{t}) \exp(i\tilde{s}s) ds dt \right\|^{p'} d\tilde{s} d\tilde{t} \\
&\quad \left(\text{because } \left| \frac{\tilde{s}/2}{\sin(\tilde{s}/2)} \right| \leq \frac{\pi}{2} \text{ for } -\pi \leq \tilde{s} \leq \pi \right) \\
&\leq \frac{1}{2\pi} \left(\frac{\pi}{2} \right)^{p'} \int_{\mathbb{G}'} \int_{\mathbb{R}} \left\| \int_{\mathbb{G}} \sum_m T \mathbf{g}_m(t) \chi_{[m-\frac{1}{2}, m+\frac{1}{2}]}(s) (t, \tilde{t}) \exp(i\tilde{s}s) ds dt \right\|^{p'} d\tilde{s} d\tilde{t} \\
&= \left(\frac{\pi}{2} \right)^{p'} \| [F^{\mathbb{R} \times \mathbb{G}}, T] \left(\sum_m \mathbf{g}_m \chi_{[m-\frac{1}{2}, m+\frac{1}{2}]} \right) \|^{p'} \\
&\leq \left(\frac{\pi}{2} \right)^{p'} \| T | \mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} \|^{p'} \left\| \sum_m \mathbf{g}_m \chi_{[m-\frac{1}{2}, m+\frac{1}{2}]} | L_p(\mathbb{R} \times \mathbb{G}) \right\|^{p'} \\
&\quad \left(\text{since } \left\| \sum_m \mathbf{g}_m \chi_{[m-\frac{1}{2}, m+\frac{1}{2}]} | L_p(\mathbb{R} \times \mathbb{G}) \right\| = \left(\int_{\mathbb{G}} \sum_m \| \mathbf{g}_m(t) \|^p dt \right)^{\frac{1}{p}} \right) \\
&= \left(\frac{\pi}{2} \right)^{p'} \| T | \mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} \|^{p'} \| \mathbf{f} | L_p(\mathbb{Z} \times \mathbb{G}) \|^{p'} \quad \text{by (2).}
\end{aligned}$$

Therefore we have $\| T | \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} \| \leq \frac{\pi}{2} \| T | \mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} \|$ and $\mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} \subseteq \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}$. This completes the proof of Proposition 1. $\square$

According to Theorem 6.3 of [4], for any LCA group $\mathbb{G}$ and operator T , $\| T | \mathcal{FT}_p^{\mathbb{G}} \| = \| T' | \mathcal{FT}_p^{\mathbb{G}'} \|$. By applying this property, we have the following:

Proposition 2. *For every LCA group $\mathbb{G}$, $\mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}}$.*

Proof. By Proposition 1 we have

$$\| T | \mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} \| = \| T' | \mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}'} \| \leq \| T' | \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}'} \| = \| T | \mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}} \|.$$

Similarly

$$\| T | \mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}} \| \leq \frac{\pi}{2} \| T | \mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} \|.$$

$\square$

From Proposition 1 and 2, we conclude that $\mathcal{FT}_p^{\mathbb{R} \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}}$ for every LCA group $\mathbb{G}$. And we have the following two corollaries.

Corollary 3. $\mathcal{FT}_p^{\mathbb{R}^n} = \mathcal{FT}_p^{\mathbb{Z}^n} = \mathcal{FT}_p^{\mathbb{T}^n}$ for every integer $n \geq 1$.

Proof. By Proposition 1 we have

$$\mathcal{FT}_p^{\mathbb{R}^n} = \mathcal{FT}_p^{\mathbb{R} \times \mathbb{R}^{n-1}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{R}^{n-1}}.$$

We continue this to obtain

$$\mathcal{FT}_p^{\mathbb{R} \times \mathbb{Z} \times \mathbb{R}^{n-2}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{Z} \times \mathbb{R}^{n-2}} = \dots = \mathcal{FT}_p^{\mathbb{Z}^n}.$$

Similarly we apply Proposition 2 n times to obtain

$$\mathcal{FT}_p^{\mathbb{R}^n} = \mathcal{FT}_p^{\mathbb{T}^n}.$$

□

Corollary 4. $\mathcal{FT}_p^{\mathbb{R}^n \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z}^n \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{T}^n \times \mathbb{G}}$ for every LCA group $\mathbb{G}$.

Proof. The proof is similar to that of Corollary 3. □

Lemma 5. For every LCA group $\mathbb{G}$ and every positive integer n ,

$$\|T|\mathcal{FT}_p^{\mathbb{Z}^n \times \mathbb{G}}\| = \|T|\mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}\|$$

and therefore $\mathcal{FT}_p^{\mathbb{Z}^n \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}$.

Proof. Let $\mathbf{f}(k, t) = \sum_m \mathbf{g}_m(t) \chi_{\{m\}}(k)$, where $\mathbf{g}_m$ is an X -valued simple function on $\mathbb{G}$ and the summation is on a finite subset of $\mathbb{Z}$, then

$$\|\mathbf{f}|L_p(\mathbb{Z} \times \mathbb{G})\| = \left(\int_{\mathbb{G}} \sum_m \|\mathbf{g}_m(t)\|^p dt \right)^{\frac{1}{p}}.$$

We have

$$[F^{\mathbb{Z} \times \mathbb{G}}, T]\mathbf{f}(\tilde{s}, \tilde{t}) = \int_{\mathbb{G}} \sum_k \left(\sum_m T\mathbf{g}_m(t) \chi_{\{m\}}(k) \right) \exp(i\tilde{s}k)(t, \tilde{t}) dt$$

and

$$\begin{aligned} \|[F^{\mathbb{Z} \times \mathbb{G}}, T]\mathbf{f}\|^{p'} &= \int_{\mathbb{G}'} \int_{\mathbb{T}} \left\| \int_{\mathbb{G}} \sum_k \left(\sum_m T\mathbf{g}_m(t) \chi_{\{m\}}(k) \right) \exp(i\tilde{s}k)(t, \tilde{t}) dt \right\|^{p'} d\tilde{s} d\tilde{t} \\ &= \int_{\mathbb{G}'} \int_{\mathbb{T}} \int_{\mathbb{T}} \left\| \int_{\mathbb{G}} \sum_{k,l} \left(\sum_m T\mathbf{g}_m(t) \chi_{\{m\}}(k) \chi_{\{0\}}(l) \right) \exp(i\tilde{s}k) \exp(i\tilde{\xi}l)(t, \tilde{t}) dt \right\|^{p'} d\tilde{s} d\tilde{\xi} d\tilde{t} \\ &\leq \|T|\mathcal{FT}_p^{\mathbb{Z}^2 \times \mathbb{G}}\|^{p'} \left(\int_{\mathbb{G}} \sum_{k,l} \left\| \sum_m \mathbf{g}_m(t) \chi_{\{m\}}(k) \chi_{\{0\}}(l) \right\|^p dt \right)^{\frac{p'}{p}} \\ &= \|T|\mathcal{FT}_p^{\mathbb{Z}^2 \times \mathbb{G}}\|^{p'} \left(\int_{\mathbb{G}} \sum_m \|\mathbf{g}_m(t)\|^p dt \right)^{\frac{p'}{p}}, \end{aligned}$$

so $\|T|\mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}\| \leq \|T|\mathcal{FT}_p^{\mathbb{Z}^2 \times \mathbb{G}}\|$.

Conversely let

$$\mathbf{f}(k_1, k_2, t) = \sum_{l_1, l_2} \mathbf{g}_{l_1, l_2}(t) \chi_{\{l_1\}}(k_1) \chi_{\{l_2\}}(k_2),$$

where the summation is on a finite subset of $\mathbb{Z} \times \mathbb{Z}$, then we have

$$(3) \quad \|\mathbf{f}|_{L_p(\mathbb{Z}^2 \times \mathbb{G})}\| = \left(\int_{\mathbb{G}} \sum_{l_1, l_2} \|\mathbf{g}_{l_1, l_2}(t)\|^p dt \right)^{\frac{1}{p}}.$$

Since $[F^{\mathbb{Z}^2 \times \mathbb{G}}, T]\mathbf{f}(\tilde{s}_1, \tilde{s}_2, \tilde{t}) = \int_{\mathbb{G}} \sum_{l_1, l_2} T\mathbf{g}_{l_1, l_2}(t) \exp(i\tilde{s}_1 l_1) \exp(i\tilde{s}_2 l_2)(t, \tilde{t}) dt$ we have

$$(4) \quad \begin{aligned} & \| [F^{\mathbb{Z}^2 \times \mathbb{G}}, T]\mathbf{f} \|^{p'} = \\ & \int_{\mathbb{G}'} \int_{\mathbb{T}} \int_{\mathbb{T}} \left\| \int_{\mathbb{G}} \sum_{l_1, l_2} T\mathbf{g}_{l_1, l_2}(t) \exp(i\tilde{s}_1 l_1) \exp(i\tilde{s}_2 l_2)(t, \tilde{t}) dt \right\|^{p'} d\tilde{s}_1 d\tilde{s}_2 d\tilde{t} \\ & \quad (\text{ we let } A = 2 \max\{l_1\} + 1) \\ & = \int_{\mathbb{G}'} \int_{\mathbb{T}} \int_{\mathbb{T}} \left\| \int_{\mathbb{G}} \sum_{l_1, l_2} T\mathbf{g}_{l_1, l_2}(t) \exp(i\tilde{s}_1 l_1) \exp(i\tilde{s}_2 A l_2)(t, \tilde{t}) dt \right\|^{p'} d\tilde{s}_1 d\tilde{s}_2 d\tilde{t} \\ & = \int_{\mathbb{G}'} \int_{\mathbb{T}} \int_{\mathbb{T}} \left\| \int_{\mathbb{G}} \sum_{l_1, l_2} T\mathbf{g}_{l_1, l_2}(t) \exp(i\tilde{s}_1 l_1) \exp(i(\tilde{s}_1 + \tilde{s}_2) A l_2)(t, \tilde{t}) dt \right\|^{p'} d\tilde{s}_1 d\tilde{s}_2 d\tilde{t} \\ & = \int_{\mathbb{G}'} \int_{\mathbb{T}} \int_{\mathbb{T}} \left\| \int_{\mathbb{G}} \sum_{l_1, l_2} T\mathbf{g}_{l_1, l_2}(t) \exp(i\tilde{s}_1 (l_1 + A l_2)) \exp(i\tilde{s}_2 A l_2)(t, \tilde{t}) dt \right\|^{p'} d\tilde{s}_1 d\tilde{s}_2 d\tilde{t} \\ & = \int_{\mathbb{T}} \int_{\mathbb{G}'} \int_{\mathbb{T}} \left\| \int_{\mathbb{G}} \sum_{l_1, l_2} T(\mathbf{g}_{l_1, l_2}(t) \exp(i\tilde{s}_2 A l_2)) \exp(i\tilde{s}_1 (l_1 + A l_2))(t, \tilde{t}) dt \right\|^{p'} d\tilde{s}_1 d\tilde{t} d\tilde{s}_2 \\ & = \int_{\mathbb{T}} \left\| [F^{\mathbb{Z} \times \mathbb{G}}, T] \left(\sum_{l_1, l_2} (\mathbf{g}_{l_1, l_2}(t) \exp(i\tilde{s}_2 A l_2)) \chi_{\{l_1 + A l_2\}} \right) \right\|^{p'} d\tilde{s}_2 \\ & \leq \int_{\mathbb{T}} \left[\| T | \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} \|^{p'} \left(\int_{\mathbb{G}} \sum_{l_1 + A l_2} \|\mathbf{g}_{l_1, l_2}(t) \exp(i\tilde{s}_2 A l_2)\|^p dt \right)^{p'/p} \right] d\tilde{s}_2 \\ & = \| T | \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} \|^{p'} \left(\int_{\mathbb{G}} \sum_{l_1 + A l_2} \|\mathbf{g}_{l_1, l_2}(t)\|^p dt \right)^{p'/p} \\ & = \| T | \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} \|^{p'} \left(\int_{\mathbb{G}} \sum_{l_1, l_2} \|\mathbf{g}_{l_1, l_2}(t)\|^p dt \right)^{p'/p} \\ & = \| T | \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} \|^{p'} \|\mathbf{f}|_{L_p(\mathbb{Z}^2 \times \mathbb{G})}\|^{p'} \quad \text{by (3),} \end{aligned}$$

so it follows that $\| T | \mathcal{FT}_p^{\mathbb{Z}^2 \times \mathbb{G}} \| \leq \| T | \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}} \|$.

Therefore we have $\|T|\mathcal{FT}_p^{\mathbb{Z}^2 \times \mathbb{G}}\| = \|T|\mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}\|$. A simple generalization yields, for all positive integers n that

$$(5) \quad \|T|\mathcal{FT}_p^{\mathbb{Z}^n \times \mathbb{G}}\| = \|T|\mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}\|$$

hence that $\mathcal{FT}_p^{\mathbb{Z}^n \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}$. $\square$

Remark 1. In fact, a dissipative group $\mathbb{A}$ satisfies the property that $\|T|\mathcal{FT}_p^{\mathbb{A}^n \times \mathbb{G}}\| = \|T|\mathcal{FT}_p^{\mathbb{A} \times \mathbb{G}}\|$ for any positive integer n and for any LCA group $\mathbb{G}$. The definition and properties of a dissipative group are introduced in [4]. The idea of the proof of Lemma 5 goes back to [4].

Corollary 6. $\|T|\mathcal{FT}_p^{\mathbb{T}^n \times \mathbb{G}}\| = \|T|\mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}}\|$ and so $\mathcal{FT}_p^{\mathbb{T}^n \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}}$.

Proof. This follows from Lemma 5 and a duality argument. $\square$

Proposition 7. Let $\mathbb{H}$ be $\mathbb{R}$, $\mathbb{Z}$ or $\mathbb{T}$, and $\mathbb{G}$ be an LCA group. Then $\mathcal{FT}_p^{\mathbb{H}^n \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}$ for any integer $n \geq 1$. In particular, $\mathcal{FT}_p^{\mathbb{H}^n} = \mathcal{FT}_p^{\mathbb{Z}}$.

Proof. It follows from Corollary 4 and Lemma 5 that $\mathcal{FT}_p^{\mathbb{H}^n \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}$. And if $\mathbb{G}$ is the trivial group then we have $\mathcal{FT}_p^{\mathbb{H}^n} = \mathcal{FT}_p^{\mathbb{Z}}$. $\square$

Notation: From now on, when $\mathbb{H}$ is $\mathbb{R}$, $\mathbb{Z}$ or $\mathbb{T}$ we denote $\mathcal{FT}_p^{\mathbb{H}}$ by $\mathcal{FT}_p$.

Corollary 8. $\mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{T}^c} = \mathcal{FT}_p$ for any nonnegative integers a, b, c with $a + b + c \geq 1$.

Proof. Without loss of generality we assume $a, b, c > 0$. By applying Proposition 7, we have

$$\mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{T}^c} = \mathcal{FT}_p^{\mathbb{Z}^{b+1} \times \mathbb{T}^c} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{T}^c} = \mathcal{FT}_p^{\mathbb{Z}^2} = \mathcal{FT}_p^{\mathbb{Z}}.$$

$\square$

Theorem 9. Let $\mathbb{G}$ be an LCA group then $\mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{T}^c \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}$ for any nonnegative integers a, b, c with $a + b + c \geq 1$.

Proof. By Proposition 7 we have

$$\mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{T}^c \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z}^{1+b} \times \mathbb{T}^c \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z}^{2+b} \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}.$$

$\square$

3. SEARCH FOR $\mathbb{G}$ SATISFYING $\mathcal{FT}_p^{\mathbb{G}} = \mathcal{FT}_p^{\mathbb{Z}}$ AND FURTHER CLASSIFICATION.

We notice that to solve the problem of deciding whether $\mathcal{FT}_p^{\mathbb{G}}$ depends on $\mathbb{G}$, we should solve that problem on the compact abelian groups since any compactly generated LCA group is a product of finite number of $\mathbb{R}$'s and $\mathbb{Z}$'s and a compact group [5]. The dual group of a compact group has the discrete topology [12]. More over if $\mathbb{G}$ is a compact abelian group then $\mathbb{G}$ is connected iff $\mathbb{G}'$ is torsion-free [5]. This fact gives a clue to the results in this section.

First, we introduce Weil's formula which factorizes an integration on $\mathbb{G}$ into double ones on a closed subgroup and its factor group.

Theorem 10. [11] *Let $\mathbb{G}$ be an LCA group and $\mathbb{H}$ a closed subgroup. Then there are Haar measures $\mu_{\mathbb{G}}$, $\mu_{\mathbb{H}}$ and $\mu_{\mathbb{G}/\mathbb{H}}$ such that*

$$(6) \quad \int_{\mathbb{G}} \mathbf{f}(s) d\mu_{\mathbb{G}}(s) = \int_{\mathbb{G}/\mathbb{H}} \left(\int_{\mathbb{H}} \mathbf{f}(s+h) d\mu_{\mathbb{H}}(h) \right) d\mu_{\mathbb{G}/\mathbb{H}}(s + \mathbb{H})$$

if $\mathbf{f}$ is a compactly supported continuous Banach space-valued function or a non-negative lower semi continuous function on $\mathbb{G}$.

In Theorem 10 if any two of $\mu_{\mathbb{G}}$, $\mu_{\mathbb{H}}$ and $\mu_{\mathbb{G}/\mathbb{H}}$ are given then the third is determined so that the statement holds.

Andersson [1] obtained the inequality, $\|I_X |\mathcal{FT}_p^{\mathbb{H}}|\| \leq \|I_X |\mathcal{FT}_p^{\mathbb{G}}|\|$, where $\mathbb{H}$ is an open subgroup of an LCA group $\mathbb{G}$ and I_X is the identity operator on a Banach space X . By replacing I_X with a bounded linear operator $T : X \rightarrow Y$ in the proof of [1], we have the following:

Proposition 11. *Let $\mathbb{H}$ be an open subgroup of an LCA group $\mathbb{G}$ then we have*

$$(7) \quad \|T |\mathcal{FT}_p^{\mathbb{H}}|\| \leq \|T |\mathcal{FT}_p^{\mathbb{G}}|\|.$$

Now we consider torsion-free LCA groups with the discrete topology.

Corollary 12. *For any non-trivial torsion-free LCA group $\mathbb{E}$ with the discrete topology,*

$$(8) \quad \|T |\mathcal{FT}_p^{\mathbb{E}}|\| = \|T |\mathcal{FT}_p^{\mathbb{Z}}|\|.$$

Proof. The proof is similar to that of $\mathcal{FT}_p^{\mathbb{E}} = \mathcal{FT}_p^{\mathbb{Z}}$ in [1]: Since $\mathbb{Z}$ is isomorphic with an open subgroup of $\mathbb{E}$, by applying (7) we have $\|T |\mathcal{FT}_p^{\mathbb{Z}}|\| \leq \|T |\mathcal{FT}_p^{\mathbb{E}}|\|$. On the other hand for any X -valued simple function $\mathbf{f}$ defined on $\mathbb{E}$ which has finite L_p -norm, $\mathbf{f}$ is non-zero only on a subset of an open subgroup which is isomorphic with $\mathbb{Z}^k$ for some positive integer k . Therefore we have

$$\frac{\|[F^{\mathbb{E}}, T]\mathbf{f}\|_{p'}}{\|\mathbf{f}\|_{L_p(\mathbb{E})}} = \frac{\|[F^{\mathbb{Z}^k}, T]\mathbf{f}\|_{p'}}{\|\mathbf{f}\|_{L_p(\mathbb{Z}^k)}} \leq \|T |\mathcal{FT}_p^{\mathbb{Z}^k}|\|$$

and hence

$$\|T |\mathcal{FT}_p^{\mathbb{E}}|\| \leq \sup_k \|T |\mathcal{FT}_p^{\mathbb{Z}^k}|\| = \|T |\mathcal{FT}_p^{\mathbb{Z}}|\|.$$

Thus the equality (8) follows. □

Lemma 13. *Let $\mathbb{H}$ be a closed subgroup of an LCA group $\mathbb{G}$ such that $\mathbb{G}/\mathbb{H}$ is finite. If n is the cardinal number of $\mathbb{G}/\mathbb{H}$ then*

$$(9) \quad \|T|\mathcal{FT}_p^{\mathbb{H}}\| \leq \|T|\mathcal{FT}_p^{\mathbb{G}}\| \leq \|T|\mathcal{FT}_p^{\mathbb{H}}\| \cdot n^{1/p'},$$

hence $\mathcal{FT}_p^{\mathbb{G}} = \mathcal{FT}_p^{\mathbb{H}}$.

Proof. Since $\mathbb{G}/\mathbb{H}$ is finite, $\mathbb{H}$ is open and by the inequality (7) we have the left side. The proof of the right side is as follows. Let us choose $s_1, s_2, \dots, s_n$ in $\mathbb{G}$ such that $\mathbb{G}/\mathbb{H} = \{s_1 + \mathbb{H}, s_2 + \mathbb{H}, \dots, s_n + \mathbb{H}\}$. The measure of $\mathbb{G}/\mathbb{H}$ is the Haar measure of unit mass. By applying Weil's formula, for any compactly supported continuous X -valued function $\mathbf{f}$ on $\mathbb{G}$ we have

$$(10) \quad \begin{aligned} \int_{\mathbb{G}} \|\mathbf{f}(s)\|^p d_{\mu_{\mathbb{G}}}(s) &= \int_{\mathbb{G}/\mathbb{H}} \left(\int_{\mathbb{H}} \|\mathbf{f}(s+h)\|^p d_{\mu_{\mathbb{H}}}(h) \right) d_{\mu_{\mathbb{G}/\mathbb{H}}}(s + \mathbb{H}) \\ &= \sum_{i=1}^n \frac{1}{n} \int_{\mathbb{H}} \|\mathbf{f}(s_i + h)\|^p d_{\mu_{\mathbb{H}}}(h) \end{aligned}$$

and

$$(11) \quad \begin{aligned} [F^{\mathbb{G}}, T]\mathbf{f}(\sigma) &= \sum_{i=1}^n \frac{1}{n} \left(\int_{\mathbb{H}} T\mathbf{f}(s_i + h) \sigma(s_i + h) d_{\mu_{\mathbb{H}}}(h) \right) \\ &= \sum_{i=1}^n \frac{\sigma(s_i)}{n} \left(\int_{\mathbb{H}} T\mathbf{f}(s_i + h) \sigma(h) d_{\mu_{\mathbb{H}}}(h) \right) \end{aligned}$$

for $\sigma \in \mathbb{G}'$. The dual group of $\mathbb{G}/\mathbb{H}$ is isomorphic with the closed subgroup $\mathbb{H}^{\perp} = \{\chi \in \mathbb{G}' \mid \chi(h) = 1 \text{ for all } h \in \mathbb{H}\}$ and $\mathbb{G}'/\mathbb{H}^{\perp}$ is isomorphic with $\mathbb{H}'$. Here the cardinal number of $\mathbb{H}^{\perp}$ is n , the measure of $\mathbb{H}^{\perp}$ is the counting measure which make Parseval's identity clean with constant 1 and we write $\mathbb{H}^{\perp}$ as $\{\tilde{\eta}_1, \tilde{\eta}_2, \dots, \tilde{\eta}_n\}$. $[F^{\mathbb{G}}, T](\mathbf{f})$ belongs to $[C_0(\mathbb{G}'), Y]$, so Y -norm of $[F^{\mathbb{G}}, T](\mathbf{f})$ is continuous and by

Weil's formula we have

$$\begin{aligned}
\| [F^{\mathbb{G}}, T] \mathbf{f} \| &= \left(\int_{\mathbb{H}'} \sum_{j=1}^n \left\| \frac{1}{n} \sum_{i=1}^n \sigma(s_i) \eta_j(s_i) \int_{\mathbb{H}} T \mathbf{f}(s_i + h) \sigma(h) d_{\mu_{\mathbb{H}}}(h) \right\|^{p'} d_{\mu_{\mathbb{H}'}}(\sigma + \mathbb{H}^{\perp}) \right)^{1/p'} \\
&\leq \left[\left(\int_{\mathbb{H}'} \sum_{j=1}^n \left(\sum_{i=1}^n \frac{1}{n} \left\| \int_{\mathbb{H}} T \mathbf{f}(s_i + h) \sigma(h) d_{\mu_{\mathbb{H}}}(h) \right\|^p \right)^{p'/p} d_{\mu_{\mathbb{H}'}}(\sigma + \mathbb{H}^{\perp}) \right)^{p/p'} \right]^{1/p} \\
&\leq \left[\sum_{i=1}^n \frac{1}{n} \left(\int_{\mathbb{H}'} \sum_{j=1}^n \left\| \int_{\mathbb{H}} T \mathbf{f}(s_i + h) \sigma(h) d_{\mu_{\mathbb{H}}}(h) \right\|^{p'} d_{\mu_{\mathbb{H}'}}(\sigma + \mathbb{H}^{\perp}) \right)^{p/p'} \right]^{1/p} \\
&= \left[\sum_{i=1}^n \frac{1}{n} n^{p/p'} \left(\int_{\mathbb{H}'} \left\| \int_{\mathbb{H}} T \mathbf{f}(s_i + h) \sigma(h) d_{\mu_{\mathbb{H}}}(h) \right\|^{p'} d_{\mu_{\mathbb{H}'}}(\sigma + \mathbb{H}^{\perp}) \right)^{p/p'} \right]^{1/p} \\
&\leq \left[\sum_{i=1}^n \frac{1}{n} n^{p/p'} \| T | \mathcal{FT}_p^{\mathbb{H}} \| ^p \left(\int_{\mathbb{H}} \| \mathbf{f}(s_i + h) \|^p d_{\mu_{\mathbb{H}}}(h) \right) \right]^{1/p} \\
&= n^{1/p'} \| T | \mathcal{FT}_p^{\mathbb{H}} \| \| \mathbf{f} \|_{L_p(\mathbb{G})}.
\end{aligned}$$

We used Minkowski's inequality in the third line. $\square$

Proposition 14. *If $\mathbb{F}$ is an infinite, compact and connected LCA group then for any nonnegative integers a, b , there exist $c(a, b), C(a, b) > 0$ such that*

$$c(a, b) \| T | \mathcal{FT}_p^{\mathbb{T}} \| \leq \| T | \mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{F}} \| \leq C(a, b) \| T | \mathcal{FT}_p^{\mathbb{T}} \|.$$

Therefore we have $\mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{F}} = \mathcal{FT}_p$.

Proof. If $a = b = 0$ then by applying (8) we have $\| T' | \mathcal{FT}_p^{\mathbb{F}'} \| = \| T' | \mathcal{FT}_p^{\mathbb{Z}} \|$ where $\mathbb{F}'$ is the dual group of $\mathbb{F}$ and hence a nontrivial torsion-free group with the discrete topology. Therefore it follows that

$$\begin{aligned}
(12) \quad & \| T | \mathcal{FT}_p^{\mathbb{F}} \| = \| T' | \mathcal{FT}_p^{\mathbb{Z}} \| = \| T | \mathcal{FT}_p^{\mathbb{T}} \| \quad \text{and} \\
& \mathcal{FT}_p^{\mathbb{F}} = \mathcal{FT}_p.
\end{aligned}$$

If $a + b \geq 1$ then by applying Theorem 9 and Corollary 4, we have

$$\mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{F}} = \mathcal{FT}_p^{\mathbb{T} \times \mathbb{F}}$$

and in fact

$$c(a, b) \| T | \mathcal{FT}_p^{\mathbb{T} \times \mathbb{F}} \| \leq \| T | \mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{F}} \| \leq C(a, b) \| T | \mathcal{FT}_p^{\mathbb{T} \times \mathbb{F}} \|$$

for some positive reals $c(a, b), C(a, b)$. Here $c(a, b) = (\frac{2}{\pi})^{ab}$, $C(a, b) = 1$. Note that $\mathbb{T} \times \mathbb{F}$ is connected and compact. Thus we again have

$$\| T | \mathcal{FT}_p^{\mathbb{T} \times \mathbb{F}} \| = \| T | \mathcal{FT}_p^{\mathbb{T}} \| \quad \text{by (12).}$$

$\square$

If $\mathbb{F}$ has only finitely many components and $\mathbb{C}$ is the component(maximal connected set) of the identity element then $\mathbb{F}/\mathbb{C}$ is finite and we have the following result.

Theorem 15. *Let $\mathbb{F}$ be an infinite compact LCA group with n components. Then for any nonnegative integers a, b , there exists $c(a, b)$, $C(a, b) > 0$ such that*

$$c(a, b) \|T| \mathcal{FT}_p^{\mathbb{T}}\| \leq \|T| \mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{F}}\| \leq n^{1/p'} C(a, b) \|T| \mathcal{FT}_p^{\mathbb{T}}\|.$$

And therefore $\mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{F}} = \mathcal{FT}_p$.

Proof. Let $\mathbb{C}$ be the component of the identity element of $\mathbb{F}$ then the factor group $(\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{F})/(\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{C})$ is a finite group with n elements. Thus by Lemma 13,

$$\|T| \mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{C}}\| \leq \|T| \mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{F}}\| \leq n^{1/p'} \|T| \mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{C}}\|$$

and

$$\mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{F}} = \mathcal{FT}_p^{\mathbb{R}^a \times \mathbb{Z}^b \times \mathbb{C}}.$$

Then the statement follows from Proposition 14. $\square$

This Theorem can not say anything about the case when $\mathbb{F}$ has infinitely many components, because of the factor $n^{1/p'}$.

Theorem 15 can be extended when our scope goes beyond the boundary of compactly generated LCA groups.

Theorem 16. *Let $\mathbb{E}$ be a nontrivial torsion free group with the discrete topology and $\mathbb{F}$ an infinite, compact and connected LCA group. Then for any LCA group $\mathbb{G}$,*

$$\|T| \mathcal{FT}_p^{\mathbb{E} \times \mathbb{G}}\| = \|T| \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}\|$$

and

$$\|T| \mathcal{FT}_p^{\mathbb{F} \times \mathbb{G}}\| = \|T| \mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}}\|.$$

If $\tilde{\mathbb{F}}$ is a compact LCA group with n components then for any nonnegative integer k there exist $c(k), C(k)$ such that

$$c(k) \|T| \mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}}\| \leq \|T| \mathcal{FT}_p^{\mathbb{R}^k \times \mathbb{E}^l \times \tilde{\mathbb{F}}^m \times \mathbb{G}}\| \leq n^{m/p'} C(k) \|T| \mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}}\|$$

and hence

$$\mathcal{FT}_p^{\mathbb{R}^k \times \mathbb{E}^l \times \tilde{\mathbb{F}}^m \times \mathbb{G}} = \mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}}$$

where $l, m = 0$ or 1 and $k + l + m \geq 1$. In particular if $\mathbb{G}$ is the trivial group then

$$\mathcal{FT}_p^{\mathbb{R}^k \times \mathbb{E}^l \times \tilde{\mathbb{F}}^m} = \mathcal{FT}_p.$$

Proof. First, $\mathbb{Z}$ is isomorphic with an open subgroup of $\mathbb{E}$, and then $\mathbb{Z} \times \mathbb{G}$ is an open subgroup of $\mathbb{E} \times \mathbb{G}$. By applying (7) we have $\|T| \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}\| \leq \|T| \mathcal{FT}_p^{\mathbb{E} \times \mathbb{G}}\|$. Conversely, for any simple X -valued function $\mathbf{f}$ which is defined on $\mathbb{E} \times \mathbb{G}$ and has finite L_p -norm, the support of $\mathbf{f}$ is a subset of $\mathbb{Z}^k \times \mathbb{G}$ for some positive integer

k . Therefore by Lemma 5 it follows that $\|T| \mathcal{FT}_p^{\mathbb{E} \times \mathbb{G}}\| \leq \sup_k \|T| \mathcal{FT}_p^{\mathbb{Z}^k \times \mathbb{G}}\| = \|T| \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}}\|$.

Second, from the duality argument it follows that

$$\|T| \mathcal{FT}_p^{\mathbb{F} \times \mathbb{G}}\| = \|T'| \mathcal{FT}_p^{\mathbb{F}' \times \mathbb{G}'}\| = \|T'| \mathcal{FT}_p^{\mathbb{Z} \times \mathbb{G}'}\| = \|T| \mathcal{FT}_p^{\mathbb{T} \times \mathbb{G}}\|.$$

The rest follows by applying the above two results, Proposition 1, 2 and Lemma 13. $\square$

Remark 2. If $\mathbb{G}$ is an LCA group and $\mathbb{C}$ is the component of the identity of $\mathbb{G}$ then $\mathbb{C}$ is a closed normal subgroup of $\mathbb{G}$ and the factor group $\mathbb{G}/\mathbb{C}$ is totally disconnected and Hausdorff. From (24.45) of [5], if $\mathbb{C}$ is open then $\mathbb{G}$ is isomorphic with $\mathbb{C} \times (\mathbb{G}/\mathbb{C})$. In particular $\mathbb{C}$ is open when $\mathbb{G}$ is locally connected. Hence to solve the question, denoted by (P), whether $\|T| \mathcal{FT}_p^{\mathbb{G}}\|$ is equivalent to $\|T| \mathcal{FT}_p^{\mathbb{T}}\|$ or not, it is useful to find $\mathbb{C}$ and $\mathbb{G}/\mathbb{C}$. Since $\mathbb{C}$ is isomorphic with $\mathbb{R}^n \times \mathbb{K}$, where n is a nonnegative integer and $\mathbb{K}$ is a compact connected group, see Theorem 9.14 of [5], the answer for (P) is affirmative when $\mathbb{G}/\mathbb{C}$ is good.

Theorem 17. *Let $\mathbb{G}$ be an LCA group with n components. Then there are positive real numbers c and C such that*

$$(13) \quad c \|T| \mathcal{FT}_p^{\mathbb{T}}\| \leq \|T| \mathcal{FT}_p^{\mathbb{G}}\| \leq n^{1/p'} C \|T| \mathcal{FT}_p^{\mathbb{T}}\|$$

and hence $\mathcal{FT}_p^{\mathbb{G}} = \mathcal{FT}_p$.

Proof. Let $\mathbb{C}$ be the component of the identity, $\mathbb{G}/\mathbb{C}$ is a finite LCA group. Hence by Lemma 13 we have $\|T| \mathcal{FT}_p^{\mathbb{C}}\| \leq \|T| \mathcal{FT}_p^{\mathbb{G}}\| \leq n^{1/p'} \|T| \mathcal{FT}_p^{\mathbb{C}}\|$. And $\mathbb{C}$ is isomorphic with $\mathbb{R}^k \times \mathbb{K}$, where k is a nonnegative integer and $\mathbb{K}$ is a compact connected group. $\mathbb{K}$ is trivial or infinite. If $\mathbb{K}$ is trivial then by applying Proposition 2 it follows that

$$c \|T| \mathcal{FT}_p^{\mathbb{T}^k}\| \leq \|T| \mathcal{FT}_p^{\mathbb{C}}\| \leq C \|T| \mathcal{FT}_p^{\mathbb{T}^k}\|$$

for some $c, C > 0$. And by (12), $\|T| \mathcal{FT}_p^{\mathbb{T}^k}\| = \|T| \mathcal{FT}_p^{\mathbb{T}}\|$. Thus we have

$$(14) \quad c \|T| \mathcal{FT}_p^{\mathbb{T}}\| \leq \|T| \mathcal{FT}_p^{\mathbb{C}}\| \leq C \|T| \mathcal{FT}_p^{\mathbb{T}}\|.$$

If $\mathbb{K}$ is infinite, we also have (14) by Proposition 14. Therefore the inequality (13) follows, and $\mathcal{FT}_p^{\mathbb{G}} = \mathcal{FT}_p$. $\square$

Theorem 18. *If $\mathbb{C}$ is the component of the identity of an LCA group $\mathbb{G}$ and $\mathbb{G}/\mathbb{C}$ is a (nontrivial torsion free group) $\times$ (finite group with cardinal n) with the discrete topology, then $\|T| \mathcal{FT}_p^{\mathbb{C}}\| \leq \|T| \mathcal{FT}_p^{\mathbb{G}}\| \leq n^{1/p'} \|T| \mathcal{FT}_p^{\mathbb{C} \times \mathbb{Z}}\|$ and $\mathcal{FT}_p^{\mathbb{G}} = \mathcal{FT}_p$.*

Proof. We have that $\mathbb{G}$ is isomorphic with $\mathbb{C} \times (\mathbb{G}/\mathbb{C})$ because $\mathbb{C}$ is open. We can apply Proposition 11 and Theorem 16 to obtain this theorem. $\square$

As for the class of Banach spaces which has Fourier type p for an infinite LCA group, we have the following:

Corollary 19. i) Under the same conditions as Theorem 16 we have

$$\mathrm{FT}_p^{\mathbb{R}^k \times \mathbb{E}^l \times \mathbb{F}^m} = \mathrm{FT}_p.$$

ii) Under the same conditions as Theorem 18,

$$\mathrm{FT}_p^{\mathbb{G}} = \mathrm{FT}_p.$$

Acknowledgements. The author would like to thank Professor Chansun Choi and a colleague Hun Hee Lee for comments.

REFERENCES

1. M. E. Andersson, *On the vector valued Hausdorff-Young inequality*, Ark. Mat. **36**(1998), no.1, 1-30.
2. J. Bourgain, *Vector-valued Hausdorff-Young inequalities and applications*, in *Geometric Aspects of Functional Analysis*, Lecture Notes in Math. 1317, p239-249, Springer-Verlag, Berlin-Heidelberg, 1988.
3. G. B. Folland, *A course in abstract harmonic analysis*, (Studies in Advanced Mathematics) CRC Press, Boca Raton, FL, 1995.
4. J. Garcia-Cuerva; K. S. Kazaryan; V. I. Kolyada; J. L. Torrea, *The Hausdorff-Young inequality with vector-valued coefficients and applications*, translation in Russian Math. Surveys **53** (1998), no. 3, 435-513.
5. E. Hewitt; K. A. Ross, *Abstract harmonic analysis*, Vol. I. second edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 115. Springer-Verlag, Berlin-New York, 1979.
6. Frank Jones, *Lebesgue Integration on Euclidean Space*, Jones and Bartlett, 1993.
7. H. König, *On the Fourier-coefficients of vector-valued functions*, Math. Nachr. **152**(1991), 215-227.
8. S. Kwapién, *Isomorphic characterizations of inner product spaces by orthogonal series with vector-valued coefficients*, Stud.Math. **44** (1972) 583-595
9. J. Peetre, *Sur la transformation de Fourier des fonctions a valeurs vectorielles*, (French) Rend. Sem. Mat. Univ. Padova **42**(1969), 15-26.
10. A. Pietsch; J. Wenzel, *Orthonormal systems and Banach space geometry*, Encyclopedia of Mathematics and its Applications **70**. Cambridge University Press, Cambridge, 1998.
11. H. Reiter, *Classical Harmonic Analysis and Locally compact Groups*, Oxford U. Press, Oxford, 1968.
12. W. Rudin, *Fourier analysis on groups*, Interscience Tracts in Pure and Applied Mathematics, No. 12 Interscience Publishers (a division of John Wiley and Sons), New York-London 1962.

DIVISION OF APPLIED MATHEMATICS, KOREA ADVANCED INSTITUTE OF SCIENCE AND TECHNOLOGY, 373-1 KUSONG-DONG, YUSONG-GU, TAEJON 305-701, REPUBLIC OF KOREA
E-mail address: ispark@amath.kaist.ac.kr